

# Prescription for Childhood Health

ASK YOUR CHILD'S DOCTOR TO PERFORM A BLOOD LEAD TEST

Name \_\_\_\_\_ Date \_\_\_\_\_

- Research published in the New England Journal of Medicine indicates that even blood lead levels below 10µg/dL can cause intellectual impairment in children. (Richard L. Canfield, Ph.D., Charles R. Henderson, Jr., M.A., Deborah A. Cory-Slechta, Ph.D., Christopher Cox, Ph.D., Todd A. Jusko, B.S., and Bruce P. Lanphear, M.D., M.P.H. (2003). Intellectual Impairment in Children with Blood Lead Concentrations below 10 µg per Deciliter. NEJM 348:1517-1526.)
- All Medicaid/Nevada Check-Up enrolled children are **required** to have a **blood test** for lead at 12 and 24 months, or before the age of 6 years if they have never been screened before.
- Lead exposure in Southern Nevada is not exclusively associated with lead-based paint. Traditional home remedies, imported candy and other food products, gold and costume jewelry, and lead-glazed earthenware used for cooking or storing food are examples of other sources of lead exposure.

**For more information, call the Childhood Lead Poisoning Prevention Program at (702) 759-1283 or go to [www.SouthernNevadaHealthDistrict.org/clppp](http://www.SouthernNevadaHealthDistrict.org/clppp)**

SOUTHERN NEVADA HEALTH DISTRICT

# Prescription for Childhood Health

ASK YOUR CHILD'S DOCTOR TO PERFORM A BLOOD LEAD TEST

Name \_\_\_\_\_ Date \_\_\_\_\_

- Research published in the New England Journal of Medicine indicates that even blood lead levels below 10µg/dL can cause intellectual impairment in children. (Richard L. Canfield, Ph.D., Charles R. Henderson, Jr., M.A., Deborah A. Cory-Slechta, Ph.D., Christopher Cox, Ph.D., Todd A. Jusko, B.S., and Bruce P. Lanphear, M.D., M.P.H. (2003). Intellectual Impairment in Children with Blood Lead Concentrations below 10 µg per Deciliter. NEJM 348:1517-1526.)
- All Medicaid/Nevada Check-Up enrolled children are **required** to have a **blood test** for lead at 12 and 24 months, or before the age of 6 years if they have never been screened before.
- Lead exposure in Southern Nevada is not exclusively associated with lead-based paint. Traditional home remedies, imported candy and other food products, gold and costume jewelry, and lead-glazed earthenware used for cooking or storing food are examples of other sources of lead exposure.

**For more information, call the Childhood Lead Poisoning Prevention Program at (702) 759-1283 or go to [www.SouthernNevadaHealthDistrict.org/clppp](http://www.SouthernNevadaHealthDistrict.org/clppp)**

SOUTHERN NEVADA HEALTH DISTRICT

# Prescription for Childhood Health

ASK YOUR CHILD'S DOCTOR TO PERFORM A BLOOD LEAD TEST

Name \_\_\_\_\_ Date \_\_\_\_\_

- Research published in the New England Journal of Medicine indicates that even blood lead levels below 10µg/dL can cause intellectual impairment in children. (Richard L. Canfield, Ph.D., Charles R. Henderson, Jr., M.A., Deborah A. Cory-Slechta, Ph.D., Christopher Cox, Ph.D., Todd A. Jusko, B.S., and Bruce P. Lanphear, M.D., M.P.H. (2003). Intellectual Impairment in Children with Blood Lead Concentrations below 10 µg per Deciliter. NEJM 348:1517-1526.)
- All Medicaid/Nevada Check-Up enrolled children are **required** to have a **blood test** for lead at 12 and 24 months, or before the age of 6 years if they have never been screened before.
- Lead exposure in Southern Nevada is not exclusively associated with lead-based paint. Traditional home remedies, imported candy and other food products, gold and costume jewelry, and lead-glazed earthenware used for cooking or storing food are examples of other sources of lead exposure.

**For more information, call the Childhood Lead Poisoning Prevention Program at (702) 759-1283 or go to [www.SouthernNevadaHealthDistrict.org/clppp](http://www.SouthernNevadaHealthDistrict.org/clppp)**

SOUTHERN NEVADA HEALTH DISTRICT

# Prescription for Childhood Health

ASK YOUR CHILD'S DOCTOR TO PERFORM A BLOOD LEAD TEST

Name \_\_\_\_\_ Date \_\_\_\_\_

- Research published in the New England Journal of Medicine indicates that even blood lead levels below 10µg/dL can cause intellectual impairment in children. (Richard L. Canfield, Ph.D., Charles R. Henderson, Jr., M.A., Deborah A. Cory-Slechta, Ph.D., Christopher Cox, Ph.D., Todd A. Jusko, B.S., and Bruce P. Lanphear, M.D., M.P.H. (2003). Intellectual Impairment in Children with Blood Lead Concentrations below 10 µg per Deciliter. NEJM 348:1517-1526.)
- All Medicaid/Nevada Check-Up enrolled children are **required** to have a **blood test** for lead at 12 and 24 months, or before the age of 6 years if they have never been screened before.
- Lead exposure in Southern Nevada is not exclusively associated with lead-based paint. Traditional home remedies, imported candy and other food products, gold and costume jewelry, and lead-glazed earthenware used for cooking or storing food are examples of other sources of lead exposure.

**For more information, call the Childhood Lead Poisoning Prevention Program at (702) 759-1283 or go to [www.SouthernNevadaHealthDistrict.org/clppp](http://www.SouthernNevadaHealthDistrict.org/clppp)**

SOUTHERN NEVADA HEALTH DISTRICT