

Southern Nevada Weekly Influenza Surveillance Snapshot CDC Week 13, March 25, 2018 to March 31, 2018

Summary:

Influenza surveillance for Clark County, Nevada includes data collected from 16 local hospital emergency rooms and 70 healthcare providers. During week 13, the influenza activity decreased in the United States and remained at the minimal level in the state of Nevada. In Clark County, as of 3/31/2018, the total number of confirmed cases of influenza during the season was 1264 with 50% being age 65 and older, and the proportion of emergency room (ER) visits for Influenza-like Illness (ILI) decreased from 4% in week 12 to 3.2% in week 13. Approximately 48% of area ER visits for ILI were children under 18 years of age. Influenza B was the dominant type circulating in Week 13. Twenty-six influenza-associated deaths, including 3 deaths of children under age 18, occurred this season in Clark County. Southern Nevada Health District will continue to update the public on the progression of the season and encourage influenza vaccination for all persons 6 months of age and older.


Table 1: Influenza Report by Age Group, Clark County, NV(CDC Week 40, 2017 to Week 13, 2018)

Age Group	Deaths	Hospitalized	Confirmed Cases*
0-4	1	43	74
05-17	2	29	99
18-24	1	29	34
25-49	1	113	181
50-64	6	172	240
65+	15	539	636
Total Confirmed Cases	26	925	1264

Table 2: Influenza Report by Virus Strain, Clark County, NV (CDC Week 40, 2017 to Week 13, 2018)

Influenza Type	Deaths	Hospitalized	Confirmed Cases*
A (2009 H1N1)	0	3	5
A (RIDT)	14	610	610
A (not sub-typed)	2	153	364
A (not sub-typed)	0	0	1
A (seasonal H3)	4	22	36
B (RIDT)	1	105	105
B (non-RIDT)	5	32	143
Total Confirmed Cases	26	925	1264

Figure 1: Weekly Count of Reported Positive Influenza Tests by Test Type**, Clark County, NV






Figure 2: Distribution of Confirmed Cases* by Virus Strain, Clark County, NV

Figure 3: Weekly Percentage of Emergency Room Visits for Influenza-Like Illness, Clark County, NV




Figure 5: Seasonal Trends of Confirmed Cases of Influenza, Clark County, NV




Figure 4: Age Distribution of Emergency Room Visits for Influenza-Like Illness, Clark County, NV




Figure 6: National ILI Activity Level- 2017-2018 Influenza Season CDC Week 13 ending March 31, 2018



*Confirmed Cases= Laboratory confirmed by PCR, culture, immnofluorescent DFA antigen staining, IHC staining, rapid influenza diagnostic tests (RIDTs) with hospitalizations for 24 hours or longer, or RIDTs with death.

**Test Type-RIDT: Rapid Influenza Diagnostic Test.

Test Type- Confirmed: PCR; Culture; Immunofluorescence, Direct (DFA) or Indirect (IFA) Florescent Antibody Staining; and Immunohistochemical (IHC)staining.

Note: The total number of confirmed cases presented in this report is subject to changes due to possible delays in reporting and processing. For additional information regarding SNHD influenza surveillance, please visit us at http://southernnevadahealthdistrict.org/stats-reports/influenza-surveillance.php.

Office of Epidemiology and Disease Surveillance (OEDS) Zuwen Qiu-Shultz, MPH, Epidemiologist