

# Southern Nevada Influenza Surveillance Update

Update Number 4: November 10, 2009

### **Summary**

For the week ending November 7, 2009:

- Influenza activity continues throughout Southern Nevada
- There is currently no evidence of increased severity of disease in Southern Nevada or the US
- Local and national laboratory surveillance indicates that nearly all reported cases of influenza are the result of 2009 Influenza A (H1N1)
- 799 cases, including 48 hospitalizations and 4 deaths were reported to the health district. The deaths include a 54-year-old male with unknown underlying health conditions, a 54-year-old female with underlying health conditions, and the first reports of pediatric deaths for 2009 influenza A (H1N1) in Clark County in a 17-year-old male and a 6-year-old male, both with underlying health conditions.

#### **Current Status**

#### Circulation

Over the first week of November, influenza activity continued both in Southern Nevada and nationwide. This is consistent with national trends, as all 10 geographic regions of the country are reporting elevated levels of influenza, and 48 states are reporting widespread geographic distribution of influenza (Source: CDC FluView). Laboratory testing both locally and nationwide is showing a stabilization or decrease in the testing positivity rate (Figure 1.1 and Table 1.1). Sentinel provider reports of patients seeking care for influenza-like illness (Figure 2.1) has displayed the same trend (Figure 2.1). The number of patients hospitalized for influenza has returned to the levels seen throughout October after a decrease in the previous week (Figure 3.4 and Table 3.1).

#### Severity

There is currently no evidence of increased severity of disease in Southern Nevada or in the United States. An indicator of the severity of disease, the proportion of hospitalized patients requiring intensive care unit admission, has decreased over the past two weeks. During September and early October, about one-third of hospitalized patients were admitted to the intensive care unit; for the week ending November 7, 23% of hospitalized patients required admission to the intensive care unit. Four

Data for the Week Ending November 7, 2009

influenza-related deaths were reported last week, including the first reports of pediatric deaths in Clark County. Three of the four deaths occurred in persons who were at higher risk for serious disease as a result of underlying health conditions (Figure 3.5); underlying health conditions were unknown for the fourth death.

## **Circulating Strains**

Local and national laboratory surveillance indicates that nearly all reported cases of influenza are the result of 2009 Influenza A (H1N1). Local pediatric laboratory surveillance has identified no seasonal influenza A H1 or H3 infections and only one influenza B infection out of 214 samples tested since the beginning of influenza season (Figure 1.1 and Table 1.1). This is consistent with national surveillance (Figure 1.2 and Table 1.2).


#### **Antiviral Resistance**

The circulating strain of 2009 Influenza A (H1N1) continues to display sensitivity to oseltamivir and zanamivir and resistance to adamantanes. Although sporadic cases of oseltamivir-resistance have been identified in the United States, nearly all patients had documented treatment or prophylaxis with oseltamivir, and occasional development of oseltamivir resistance during treatment or prophylaxis is not unexpected. Since April of 2009, a total of 14 cases of oseltamivir resistance have been identified in the United States. Twelve of these patients had documented exposure to oseltamivir through either treatment or chemoprophylaxis, one patient had no documented oseltamivir exposure, and one patient is under investigation to determine exposure to oseltamivir. (Source: CDC - http://www.cdc.gov/flu/ weekly/).

# Section One: Laboratory Surveillance

Enhanced pediatric influenza surveillance (EPIS) is conducted through four Clark County, NV medical practices. Each practice submits up to 10 specimens each week from pediatric patients presenting with respiratory disease and the specimens are tested for influenza and typed by RT-PCR. National surveillance is conducted through laboratories participating in the Center for Disease Control and Prevention (CDC) National Respiratory and Enteric Virus Surveillance System (NREVSS) program.

Figure 1.1 Proportion of Influenza Viruses - Clark County Pediatric Influenza Surveillance



Note: The category of "seasonal flu" includes all influenza A types other than 2009 H1N1 and all influenza B. Source: EPIS

Table 1.1 Laboratory Testing - Clark County Pediatric Influenza Surveillance

			Season to Date									
	10	10/10		10/17		10/24		10/31		_/7	From 8/30/09	
Testing Category	n	%	n	%	n	%	n	%	n	%	n	%
Influenza Negative	9	31	7	23	9	36	11	46	11	69	99	43
2009 H1N1 Positive	20	69	24	77	16	64	13	54	5	31	128	56
Flu A H1 (seasonal) Positive	0	0	0	0	0	0	0	0	0	0	0	0
Flu A H3 (seasonal) Positive	0	0	0	0	0	0	0	0	0	0	0	0
Flu B Positive	0	0	0	0	0	0	0	0	0	0	1	0
Specimens Tested	29		31		25		24		16		228	

Source: EPIS

Figure 1.2. Proportion of Influenza Viruses - National Laboratory Influenza Surveillance



Note: The category of "seasonal flu" includes all influenza A types other than 2009 H1N1 and all influenza B. Source: CDC/NRVESS

Table 1.2 Laboratory Testing Results - National Influenza Surveillance


			Season to Date							
	10/1	10/10		10/17		10/24		1	From 8/30/09	
Testing Category	n	%	n	%	n	%	n	%	n	%
Influenza Negative	9,828	71	8,088	62	11,374	58	8,893	63	71,039	69
2009 H1N1 Positive	2,505	18	3,378	26	5,453	28	3,889	27	20,669	20
Flu A H1 (seasonal) Positive	0	0 0		0	0	0	2	0	24	0
Flu A H3 (seasonal) Positive	0	0	0	0	0	0	2	0	18	0
Flu A Positive, Untyped	1,573	11	1,466	11	2,794	14	1,351	10	11,563	11
Flu B Positive	15	0	11	0	21	0	14	0	83	0
Specimens Tested	13,921		12,943		19,642		14,151		103,396	

Note: National data lags local data by one week, thus national data for the most recent week are unavailable. Source: CDC/NRVESS

#### Section Two: Sentinel Physician Influenza-Like Illness Surveillance

Data from physicians enrolled in the Center for Disease Control and Prevention's Outpatient Influenza-like Illness Surveillance Network (ILINet) indicate the percentage of all patients in a given week presenting with influenza-like illness (ILI), which is defined as a fever and either a cough or sore throat.

Figure 2.1 Percentage of Visits for Influenza-Like Illness Reported to ILINet, Locally and Nationally




Note: ILI data collection for the previous week is not complete when this report is generated, and results will lag other parts of this report by one week. Source: CDC/ILINet

#### Section Three: Clark County Reportable Disease Surveillance


Per Nevada Administrative Code 441A.575, healthcare providers and laboratories must report all laboratory-confirmed cases of influenza to the health authority. Reported hospitalizations are further investigated for the presence of underlying risk factors and for the severity of illness, including intensive care unit (ICU) admission.

Figure 3.1 Clark County Reported Influenza Hospitalization Rates by Age, Season to Date




Source: Southern Nevada Health District

Figure 3.3 Clark County Reported Influenza Deaths by Age, Season to Date



Source: Southern Nevada Health District

Figure 3.2 Clark County Reported Influenza Case Rates by Age, Season to Date




Source: Southern Nevada Health District

# Textbox 3.1 Details of Influenza-Related Deaths, Clark County, Week Ending November 07, 2009

4 deaths:

6M with underlying conditions 17M with underlying conditions 54M with underlying conditions 54M with unknown underlying conditions

Figure 3.4 Influenza Hospitalizations, Season to Date (From August 30, 2009), Clark County



Note: Data are presented by "event date", the earliest known date for a case. Although this is ideally a disease onset date, a standardized, hierarchical process is used to assign this date when the onset date is unavailable. Hospitalization that occurred before Aug. 30 but were reported Aug. 30 or after are not included in this figure, as data are displayed by the date of hospitalization.

Figure 3.5 Influenza Deaths, Season to Date (From August 30, 2009) by Date of Death, Clark County




Table 3.1 Counts Influenza Cases by Type, Hospitalizations by Type, and Deaths, Most Recent Week and Season to Date (From August 30, 2009)

Season to Da		ported \				r 07, 20	Season To Date (From August 30, 2009)							
	Cas	es Reported			Hospitalizations			Cases Reported				Hospitalizations		
Age Group	Influenza	Novel N1H1 Influenza	Total	Deaths	Non-ICU Admissions	ICU Admissions	Total Admissions	Influenza	Novel N1H1 Influenza	Total	Deaths	Non-ICU Admissions	ICU Admissions	Total Admissions
0 to 4	60	101	161	0	3	2	5	252	372	624	0	25	7	32
5 to 9	36	120	156	1	1	0	1	245	483	728	1	7	2	9
10 to 14	18	84	102	0	1	0	1	227	486	713	0	5	2	7
15 to 19	11	55	66	1	1	0	1	123	306	429	1	12	3	15
20 to 24	9	42	51	0	6	0	6	50	163	213	1	20	4	24
25 to 29	12	25	37	0	3	0	3	46	107	153	1	12	5	17
30 to 34	9	43	52	0	3	0	3	37	123	160	1	14	4	18
35 to 39	5	32	37	0	0	1	1	40	112	152	0	10	2	12
40 to 44	4	33	37	0	4	0	4	19	112	131	4	11	8	19
45 to 49	4	28	32	0	3	0	3	29	93	122	1	18	6	24
50 to 54	2	25	27	2	7	3	10	14	86	100	4	16	11	27
55 to 59	3	18	21	0	4	0	4	16	64	80	0	17	3	20
60 to 64	0	7	7	0	1	1	2	13	27	40	2	7	7	14
65 to 69	2	4	6	0	0	2	2	7	16	23	0	7	4	11
70 to 74	1	3	4	0	0	0	0	4	7	11	0	4	1	5
75 to 79	0	1	1	0	0	2	2	2	5	7	0	2	2	4
80 to 84	0	0	0	0	0	0	0	1	0	1	0	0	0	0
85+	0	2	2	0	0	0	0	1	5	6	0	0	0	0
Total	176	623	799	4	37	11	48	1,126	2,567	3,693	16	187	71	258

Note: Case and hospitalization data for the most recent week are limited to those cases reported in the one-week period ending on the date listed, and are based solely on the date in which the case was reported to SNHD. Cases listed as "Novel H1N1 Influenza" are confirmed by RT-PCR. Cases listed as "Influenza" include all patients who tested positive by a rapid influenza test and have either had no confirmatory testing or confirmatory testing indicating the presence of seasonal influenza. Case categories are mutually exclusive, as are hospitalization categories. Deaths listed are by the date or which the patient died. Cumulative totals may not add up to the current week total plus the cumulative total from the previous week, as cases from previous weeks can be reclassified from Influenza to Novel H1N1 based on new lab results, and duplicates are identified and removed.